

From: [David Hoffman](#)
To: [MN_PUC_PublicAdvisor](#)
Subject: Please file to docket #22-132 as a petition for reconsideration to the decision made on April 10th. Please see attached letter and enclosures.
Date: Wednesday, May 28, 2025 9:32:26 AM

You don't often get email from dlajhoff60@outlook.com. [Learn why this is important](#)

This message may be from an external email source.

Do not select links or open attachments unless verified. Report all suspicious emails to Minnesota IT Services Security Operations Center.

From: David Hoffman <dlajhoff60@outlook.com>
Sent: Wednesday, May 21, 2025 9:56 AM
To: David Hoffman <dlajhoff60@outlook.com>
Subject: Xcel

To the Minnesota Public Utility Commissions Board:

This letter is in regards to the Xcel transmission line route that you chose last fall in Gales/Johnsonville townships in Redwood County, MN, docket #22-132. I am of the understanding per Andrew Welch with Xcel that we have 20 days to oppose this route once the papers are officially signed.

My husband along with one of our neighbors, who lives on the proposed route, attended a meeting last fall in Marshall, MN, at the 5 Family Ranch. They were informed at this meeting that if anyone had an alternative route to the route segments 219 and 211 to mail that information in by Nov. 25th. We sent a letter addressed to you five commissioners on Nov. 21st via certified mail. You received this letter prior to the deadline. This letter outlined the pros and cons of your proposed route, and the alternative route we chose along with a petition that was signed by all the residents affected by the route you chose who were against it. We sent this letter via certified mail and it was received on November 23rd which was before the deadline that was provided to them by the Xcel representative at the meeting. I was informed recently by Andrew Welch from Xcel that none of our information was even considered or looked at by any of you because "they were already past that part of the process". So why were we given false information from the representative at the meeting? Perhaps, because they didn't want to deal with all the opposition they knew would come up at this meeting. A lot of

time and effort was spent gathering those details and making sure it was sent off in a timely fashion.

The route that you have chosen for these high voltage transmission lines has four homes with very close proximity to the highway and have a total of 9 small children, all under the age of 6. The route we proposed has NO children and the majority of the homes are further away from the highway with longer driveways. Your #1 concern when choosing these routes should be first and foremost the residents who live along the route. I have included studies that have been performed regarding the harmful effects of high voltage transmission lines on humans, animals and plants. I'm not going to insult your intelligence, but if you have no idea as to the harmful effects of HVT lines then please read the enclosed material and educate yourselves. After you have read the material, ask yourself if you would want these monstrosity lines out your front door or the door of your small children or grandchildren...I think not.

Following is a list of the pros and cons to the route you chose and the alternative route we provided last November:

PROs of moving the transmission line North to County Road 55/210th Street:

- Along this route there are NO children and there would only be 2 residences within variance distance
- It is a shorter route with 1 mile LESS transmission line
- 2.5 miles LESS of existing power line competing for right of way
- This is a well-maintained county gravel road for easy maintenance, with very low traffic (35 vehicles daily in 2019 according to MNDot traffic mapping website)
- Goes past LESS wetland areas and much LESS wildlife areas
- Crosses the Cottonwood River ONCE instead of twice
- Crosses 2 dredge ditches with NO wetlands surrounding them

CONS of transmission line staying South on Route segments 219 and 211:

- South route contains at least FOUR residences within variance distance, and NINE small children under the age of 6
- Transmission line heads in the wrong direction
- 5.5 miles of existing power line competing for right of way
- This is a highly traveled paved road, 150 vehicles a day in 2019 according to MNDot traffic mapping website
- Follows the Cottonwood River and many wetlands and crosses the river TWICE vs once on the proposed route
- Crosses 2 streams with surrounding wetlands/valleys

So, my question to all of you is WHO really chose this route? As you can see with the above pros and cons, the route we proposed is clearly the safer, cheaper route. Mr. Bob Zwach sold you the property near Garvin for the substation. He made the comment, "I don't care where they put the transmission lines as long as they don't put them on my property"! Don't penalize the residents with several small children because of a decision that was made by another who capitalized from selling Xcel that land, which I'm pretty sure he received \$20,000+ an acre. He can deal with these consequences.

You will have to use eminent domain to obtain access to every farmer's land along the route you have chosen, as clearly according to our petition everyone who owns property along that route agrees that it should be moved North to County Road 55/210th Street due to the proximity of the homes and the cost.

If the route goes down County Road 4, I can guarantee all of you that I will be monitoring the health of ALL the individuals who live along that route, especially the small children (3 of which are my grandchildren). If it affects the health of ANYONE, Xcel will have the biggest lawsuit on their hands that they have ever witnessed! I worked in the medical field for 35 years and have a wealth of knowledge into illnesses and the root causes.

I was also told that none of you actually made the trip down here to lay your eyes on the 2 routes, but yet you all made the decision on the County Road 4 route.

Thank you for your time and consideration in this very important matter. It should have been reviewed back in November. I look forward to hearing from you.

Respectfully submitted,


Lisa Hoffman

Enclosures - petition, alternate route

210th St

200th St

County Highway 4



County Highway 8

County Highway 5

Redwood County
Gales & Johnsonville Townships

— = Improved
Blue
Route
Segment

Papa's

miles North along 210th Street in Redwood county through Gales and Johnsonville townships.

Name	Address	Contact Info	Signature	Date
Clay Alan Churchill	13218 Co. Hwy. 4 Walnut Grove, MN 56180	(507) 381-6826	Clay Churchill	11/16/24
Abby Jo Churchill	13218 Co Hwy 4 Walnut Grove, MN 56180	507-829-4769	Abby Churchill	11/16/24
Brian David Hicks	19465 Co Hwy 8 Tracy, MN 56175	507-829-6405	Brian Hicks	11/16/24
Michelle Mary Hicks	19465 Co Hwy 8 Tracy, MN 56175	507-620-0701	Michelle Hicks	11/16/24
Allen Leon Rydin	14072 Co. Hwy 4 Walnut Grove ^{MN} 56180	612-269-9294	Allen Rydin	11-16-24
LINDA KAY RYDIN	14072 County Hwy 4, Walnut Grove, ^{MN} 56180	507-626-5501	Linda Rydin	11-16-24
David Penner Hoffman	13866 170 th St Walnut Grove Mn 56180	DLajhoff@outlook.com 5076401617	David Hoffman	11-16-24
LISA Ann Hoffman	13866 170 th St. Walnut Grove, MN 56180	507-626-0151	Lisa Hoffman	11-16-24
Maddylene Gail Aarsvold	19568 Camp Ave Walnut Grove, MN 56180	507-828-9121	Maddylene Aarsvold	11-16-24
Late Anthony Aarsvold	19568 Camp Ave Walnut Grove, MN 56180	507-951-0177	Late Anthony Aarsvold	11-16-24
Dennis Robert Rasmussen	17698 County Hwy 4 Walnut Grove Mn 56180	507-829-4049	Dennis Rasmussen	Nov 16 24
Andrew David Rasmussen	19565 Duncan Ave, Walnut Grove MN 56180	507-747-3211	Andrew Rasmussen	11-16-24
Rebecca Ann Rasmussen	19565 Duncan Ave, Walnut Grove, MN 56180	507-227-3413	Rebecca Rasmussen	11-16-24
James Robert Zeng	19563 Camp Ave Walnut Grove, MN 56180	507-626-1104	James Zeng	11/17/24
Rachel Ann Zeng	19563 Camp Ave Walnut Grove, MN 56180	218-640-1615	Rachel Zeng	11/17/24

...the signing this petition either live near to or own property on Highway 4 in Redwood county through Gales and Johnsonville townships. Signees vastly disapprove of proposed route and proposed road widening project. I am a resident of 11 miles North along 210th Street in Redwood county through Gales and Johnsonville townships.

Name	Address	Contact Info	Signature	Date
Zachary William Otto	17603 county Highway 4 Walnut Grove MN 56180	507-266-6066	<i>Zachary William Otto</i>	11/17/24
Rebecca Rose Otto	17603 county Hwy 4 Walnut Grove MN 56180	507-828-1990	<i>Rebecca Rose Otto</i>	11/17/24
Jane Marie Hoffman	17480 Crown Ave Walnut Grove MN 56180	507/828-9936	<i>Jane Marie Hoffman</i>	11/17/24
Jane Marie Hoffman	17480 Crown Ave Walnut Grove, MN 56180	507/828/8021	<i>Jane Marie Hoffman</i>	11/17/24
Victoria Lynn Romm	(Dallenbach) 116 Pleasant View Road, Slayton, MN 56172	507-530-2701	<i>Victoria Lynn Romm</i>	11/17/24
Victoria Lynn Romm	(Dallenbach) 901 Country Circle Dr. Tracy MN 56175	507-828-7947	<i>Victoria Lynn Romm</i>	11/17/24
Catherine Juanita Dallenbach	901 Country Circle Dr. Tracy MN 56175	507-828-4763	<i>Catherine Juanita Dallenbach</i>	11/17/24
CHAD DAVID DALLENBACH	8 Sandbar Road Tracy, MN 56175	(507)829-7348	<i>Chad Dallenbach</i>	11/18/24
Cynthia JoAnn Dallenbach	8 Sandbar Road Tracy, MN 56175	(507)828-7154	<i>Cynthia JoAnn Dallenbach</i>	11/18/24
PATRICK JOHN SCHOTT	119 BURR OAK ROAD PO BOX REDWOOD FALLS MN 56283	507-830-0610	<i>Patrick John Schott</i>	11-18-2024
Chase Chad Dallenbach	21976 Co Hwy 5 Walnut Grove MN 56180	507-828-1487	<i>Chase Chad Dallenbach</i>	11-18-24
Monica Paige Dallenbach	21976 County Highway 5 Walnut Grove, MN 56180	320-291-2177	<i>Monica Paige Dallenbach</i>	11-18-24
Chelsie Kay Lundberg	19743 Co Hwy 5 Walnut Grove, MN 56180	507-829-1671	<i>Chelsie Kay Lundberg</i>	11-18-24
Brenden Donald Lundberg	19743 County Hwy 5 Walnut Grove, MN 56180	507-430-8741	<i>Brenden Donald Lundberg</i>	11-18-24
Charlene Lynn Hennrich	16345 Bunker Ave., MN 56180	507-829-3377	<i>Charlene Lynn Hennrich</i>	11-18-24

Papa's

Research Paper



The Effect of Electrical Fields From High-voltage Transmission Line on Cognitive, Biological, and Anatomical Changes in Male Rhesus macaque Monkeys Using MRI: A Case Report Study

Hamed Aliyari¹, Hedayat Sahraei², Sahar Golabi³, Mohammad Bagher Menhaj⁴, Masoomeh Kazemi², Seyed Hossein Hosseini^{4*}

1. Department of Psychology, Center for Depression Research and Clinical Care, UT Southwestern Medical Center, Dallas, Texas, United States

2. Neuroscience Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

3. Department of Medical Physiology, School of Medicine, Abadan University of Medical Sciences, Abadan, Iran.

4. Department of Electrical Engineering, Amirkabir University of Technology, Tehran, Iran.



Citation Aliyari, H., Sahraei, H., Gholabi, S., Menhaj, M. B., Kazemi, M., and Hosseini, S. H. (2022). The Effect of Electrical Fields From High-voltage Transmission Line on Cognitive, Biological, and Anatomical Changes in Male Rhesus macaque Monkeys Using MRI: A Case Report Study. *Basic and Clinical Neuroscience*, 13(4), 433-442. <http://dx.doi.org/10.32598/bcn.2021.1340.3>

doi <http://dx.doi.org/10.32598/bcn.2021.1340.3>



Article info:

Received: 10 Jul 2021

First Revision: 09 Dec 2020

Accepted: 25 Dec 2020

Available Online: 01 Jul 2022

Keywords:

MRI, Rhesus macaque, Electromagnetic fields (EMFs), High-voltage transmission line, Adrenaline hormone

ABSTRACT

Introduction: Living near high-voltage power lines and exposure to high-frequency electromagnetic fields (EMFs) is a potentially serious hazard to animal and human health. The present study was conducted to evaluate the effect of high-frequency EMFs from simulated high-voltage electric towers on cognitive, anatomical, and biological changes in the male macaque.

Methods: In this study, two Rhesus macaque were recruited, one experimental and one control. The experimental subject was exposed to EMFs from 3 kV/m simulated electric towers with a specific protocol and the control subject was tested without irradiation (4h per day, for 30 days). All required tests were performed before and after the intervention on experimental and control monkeys. The anatomical alternation of the prefrontal area (PFA) was measured by MRI images. All tests were performed on irradiated and control animals before and after the intervention and the results were compared between irradiated and control animals.

Results: The results of the present study indicated increased white blood cell counts after high-frequency EMFs irradiation. Also, the red blood cell counts showed a decreasing trend after irradiation. The plasma adrenaline level increased after irradiation. Besides, the blood glucose levels increased after irradiation. The PFA was different before and after the irradiation. Moreover, some behavioral disorders, such as fatigue, drowsiness, anorexia, and insomnia were observed after irradiation.

Conclusion: The results of biological tests and MRI showed an elevated risk of immunodeficiency disorders, weakness, and behavioral disorders. People who live or work near high-voltage electric towers with high-frequency EMFs are warned.

* Corresponding Author:

Seyed Hossein Hosseini, PhD.

Address: Department of Electrical Engineering, Amirkabir University of Technology, Tehran, Iran.

Tel: +98 (21) 64543343

E-mail: hosseini@aut.ac.ir

Highlights

- Magnetic, and electric fields from high pressure towers caused negative effects in terms of biology and even anatomical changes in the prefrontal part of the brain.
- Disturbance in the prefrontal part of the brain caused the monkey's cognitive and behavioral disorder.
- An increase in white blood cells, a decrease in red blood cells, and an increase in the adrenaline and blood sugar were indicative of biological disorders after wave exposure in male rhesus monkeys.
- The effects of magnetic and electric fields resulting from high pressure towers on the nerves and psyche require health researchers to do more studies.

Plain Language Summary

Today, one of the factors that threaten the cognitive and behavioral health of humans and animals is living in the vicinity of magnetic and electric fields resulting from the power transmission of high-pressure towers. These fields cause cognitive and behavioral disorders in living beings. Therefore, because it threatens the cognitive health of creatures, it needs more research.

1. Introduction

Nowadays, exposure to high-frequency electromagnetic fields (EMFs) is a potentially serious hazard to animal and human health (Hamed Aliyari, Hedayat Sahraei, Mohammad Reza Daliri, Beh Minaei-Bidgoli, et al., 2018; LaBar & Cabeza, 2006). According to the literature, people exposed to EMFs are more likely to develop any type of blood cancer, abortion due to poor fetal implantation, and neurological disorders, such as stress, depression, and insomnia due to degradation in neuronal function and dysfunction by reactive oxygen species (ROS) molecule (Bayr, 2005; Garip & Akan, 2010). The ROS may harm the function of hippocampus neurons and is a major contributor to mental illnesses, such as Alzheimer's disease (Aliyari, Hosseinian, Sahraei, & Menhaj, 2018; Harach et al., 2015; Sobel & Davanipour, 1996). Electric fields and currents exist in the body of all living organisms acting as a contributor to physiological control mechanisms, such as the nervous system, cell death, growth, tissue growth, and repair (Cook, Thomas, & Prato, 2002; Paz-Caballero, Cuetos, & Dobarro, 2006). The EMFs are associated with many symptoms, such as dizziness, tinnitus, weakness and fatigue, blurred vision, and drowsiness during work, as well as the emergence of unknown diseases, altered blood composition, impaired musculoskeletal system, and immune system dysfunction among animals and humans (Kazemi et al., 2018a; McGaugh & Roozendaal, 2002; Orban et al., 2003; Elaheh Tekieh et al., 2017). Impaired secretion of hormones may lead to behavioral and cognitive disorders (Hamed

Aliyari, Hedayat Sahraei, Mohammad Reza Daliri, Beh Minaei-Bidgoli, et al., 2018). Accordingly, EMFs adversely affect the secretion of protein and amine hormones by activating G-protein coupled receptors and membrane enzymes (Kazemi et al., 2018; Elaheh Tekieh et al., 2017). The adrenocorticotrophic hormone (ACTH) secreted from the anterior pituitary, acts to stimulate the adrenal cortex hormones and regulates the secretion of cortisol and adrenaline (Fereidouni, Vahidi, Shishehgar, Mehr, & Tahmasbi, 2014; Kazemi et al., 2018). The results of many studies indicate the themes of simulated high voltage towers, the reduction of melatonin secretion resulting in sleep, attention, learning, and memory disturbances, and the development of mental illness, such as depression in Rhesus monkey (H Aliyari et al., 2018; Aliyari, Hosseinian, Menhaj, & Sahraei, 2018; Aliyari et al., 2015; Aliyari, Sahraei, Daliri, Beh Minaei-Bidgoli, et al., 2018; Aliyari, Sahraei, Erfani, Mohammadi, Kazemi, Daliri, Minaei-Bidgol, et al., 2020; Aliyari, Sahraei, Golabi, et al.; Aliyaria et al., 2019; Brady, Konkle, & Alvarez, 2011; Fereidouni et al., 2014; Kazemi et al., 2018, 2018b; Elaheh Tekieh et al., 2017).

Increased secretion of adrenaline increases glucose release from the liver leading to increased blood glucose levels. Thus, the required energy for muscle and nerves is supplied (Aliyari et al., 2015; Brady et al., 2011; Harach, 2010; Elaheh Tekieh et al., 2017). Secretion of adrenaline increases due to stress and abnormal conditions. Producing more adrenaline, in turn, helps increase the amount of energy to adapt to new conditions. Besides, further production of adrenalin may deplete sugar supplies (De Pittà, Volman, Berry, & Ben-Jacob, 2011; Hickie &

Rogers, 2011). In response to environmental factors in stress conditions, the production of T-cell lymphocytes increases (R. Liburdy, Sloma, Sokolic, & Yaswen, 1993; Liburdy, 1992; Poiesz et al., 1980). The number of lymphocytes is 70% in apes and 30% in human blood. Environmental factors and stressors can increase white blood cells (WBC) or leukocytes. The abnormal increase in WBC counts may cause immune deficiency, cognitive impairment, infectious diseases, bone marrow failure, and leukemia (Monti et al., 1991). Several studies have shown that the EMFs generated from the simulated high voltage towers reduce the area of the hippocampus and the amygdala in Rhesus monkeys (Duan et al., 2014; Elahe Tekieh et al., 2018). Therefore, this study aimed to investigate the effect of high-frequency EMFs from simulated high-voltage electric towers on cognitive, anatomical, biological, biochemical, and hormonal changes in male Rhesus macaque monkeys.

2. Materials and Methods

Animals

Two Rhesus monkeys (*Macaca mulatta*) aged 4-5 years with an average weight of 4 kg were recruited. All ethical standards regarding animal confinement, transport, location, and maintenance were observed under the international laws and regulations (IR.BUMS.REC.1394.112).

Experimentation protocol and simulation of high-frequency electromagnetic fields (EMFs)

First, pretests were conducted on study subjects. Of the two selected monkeys, one subject was irradiated by 3 kV/m high-voltage EMF, 4h per day for 30 days (Aliyari, Hosseinian, Sahraei, & Menhaj, 2019; Kazemi et al., 2018, 2018b), and one subject was maintained in a non-irradiated environment and tested without irradiation (Aliyari, Hosseinian, Sahraei, & Menhaj, 2019; Kazemi et al., 2018, 2018b). The monkeys were kept in Teflon cages which, in addition to having special conditions for keeping the animal, such as strength, portability, and washability, had special properties to conduct scientific experiments (with the highest transfer properties and lowest resistance) (H. Aliyari et al., 2019; Kazemi et al., 2018). High voltage EMF (3 kV/m) was produced and simulated by a team of experts at Amir Kabir University, Tehran City, Iran. Two metal plates (2×2m) were placed in the upper part and one in the lower part of the cage at a 2 m distance, and a 6 kV EMF was released to produce a uniform EMF of 3 kV inside the cage (1×1×1m) (H Aliyari et al., 2018; Kazemi et al., 2018a; Tekieh et al., 2017).

Behavioral tests

In this study, the monkeys were vaccinated against tetanus, pertussis, diphtheria, and polio before entering the laboratory. After entering the laboratory, it took three months to adapt to the environment and adapt to the researchers. After this period, two perfectly healthy monkeys were included in the study. The monkey that was subjected to high-pressure surges was the freshest, smartest, and most active monkey in the laboratory. For this purpose, all observable behavior in subjects, such as sleeping patterns, sedentary and natural appetite, drowsiness, and depression were observed and recorded before, during, and after intervention by video cameras (Hamed Aliyari, 2018; Aliyari, Sahraei, Erfani, et al.; Aliyari, Sahraei, Erfani, Mohammadi, Kazemi, Daliri, Minaei-Bidgoli, et al., 2020; Aliyaria et al., 2019).

Biological test

Biological tests were performed on the irradiated and control subjects before and after the intervention and during the recovery period, and the obtained data were analyzed. According to the experimental protocol, a fasting period of 15h was required for the monkeys. For biological analysis, a total of 10 cc of blood was taken in three stages of pre-, post-irradiated, and recovery from the femoral artery of the animals, of which 5 cc was used to evaluate changes in adrenaline levels using the primate custom positive selection kit (MyBioSource, USA) at three stages. Another 5 cc of blood samples were used to measure fasting blood sugar, WBC counts, and red blood cell (RBC) count in three stages (H Aliyari, 2018; Hamed Aliyari, Seyed Hossein Hosseinian, et al., 2018; Hamed Aliyari, Hedayat Sahraei, Mohammad Reza Daliri, 2018; Kazemi et al., 2018a; Elahe Tekieh et al., 2018).

The anatomical alternation of the profunda femoris artery (PFA) was measured by MRI images and digital imaging and communications in medicine (DICOM) LiteBox files before and after the intervention (Hamed Aliyari, 2018; Servien, Viskontas, Giuffre, Coolican, & Parker, 2008).

Recovery phase

Both two selected monkeys, one subject exposed to irradiation and one subject maintained in a non-irradiated environment, stayed in the same previous place and situation but both were without irradiation for up to one month (H Aliyari et al., 2019).




Figure 1. Adrenaline plasma levels (pre-test, post-test, recovery)

NEURSCIENCE

The serum adrenaline levels were shown to elevate after irradiation. The adrenaline levels during the recovery phase declined to the pre-irradiation phase.

3. Results

The behavioral results of the study showed that in the tested monkey exposed to the electric field of the simulated high-pressure rig, the behavioral-cognitive indices change over time. From the first day to the 30th day of the waving, cognitive changes were visible to the naked eye (recording behavior by camcorders). Our observations of monkeys under a high-pressure electric field showed states of impatience, lethargy, anorexia, sleep disturbance, inactivity, and state of depression.

Behavioral tests investigated observable cognitive changes in the study subjects, including drowsiness, loss of appetite, sleep deprivation, fatigue, inactivity, and depression (recorded by video cameras). Behavioral results showed an increasing trend in the men-

tioned criteria during thirty days of irradiation. The behavioral analysis was performed one month before, during, and after irradiation (recovery phase). Behavioral tests were also performed on the control animal at three stages without irradiation, and no difference was observed in the control monkey.

The serum adrenaline levels were shown to elevate after high-frequency EMFs irradiation compared to the control monkeys. In addition, the adrenaline levels declined during the recovery phase to the pre-irradiation phase. No difference was observed in the three phases in the control monkeys (Figure 1).

The fast blood glucose levels were shown to increase after high-frequency EMFs irradiation compared to the control monkeys. Also, the blood glucose levels declined




Figure 2. Fast blood sugar levels (pre-test, post-test, recovery)

NEURSCIENCE

The increasing trend of fast blood glucose levels after irradiation. The fast blood glucose levels during the recovery phase were declining to the pre-irradiation phase




Figure 3. White Blood Cells (WBC) counts (pre-test, post-test, recovery)

NEURSCIENCE

The increasing trend of WBC counts after irradiation. The WBC counts during the recovery phase declined to the pre-irradiation phase.

during the recovery phase to the pre-irradiation phase. No difference was observed in the three phases in the control monkeys (Figure 2).

The WBC counts showed an increasing trend after high-frequency EMF irradiation compared to the control monkey (Figure 3).

The RBC counts were shown to decrease after high-frequency EMFs irradiation compared to the control monkeys. Besides, the RBC counts increased during the recovery phase to the pre-irradiation phase. No difference was observed in the three phases in the control monkeys (Figure 4).

The PFA decreased after irradiation in the experimental monkey compared to the control monkeys. Also, the PFA during the recovery phase increased to the pre-irradiation phase.

diation phase. No anatomical difference was observed in the control monkey (Figures 5).

4. Discussion

Today, industrialization has remade a major part of people's life. Rapidly growing industrialized cities increase the growth of power lines and the proximity of overhead electric towers to residential and occupational places increases the risk of EMFs harm as a global public concern. According to the literature, the EMFs generated by high-pressure electric towers are a serious threat to the environment of animals and humans. This destructive environmental factor adversely affects health and lifestyle and causes cognitive and behavioral changes in animals and humans (Aliyari et al., 2015; Hamed Aliyari, Hedayat Sahraei, Mohammad Reza Daliri, Beh Minaei-Bidgoli, et al., 2018; Kazemi et al., 2018; Liburdy et al., 1993; Monti et al., 1991).




Figure 4. Red Blood Cell (RBC) counts (pre-test, post-test, recovery)

NEURSCIENCE

The RBC counts were shown to decrease after irradiation. The RBC counts during the recovery phase elevated to the pre-irradiation phase.




Figure 5. Pre-frontal Area (PFA)

NEURSCIENCE

The pre-frontal area was shown to decrease after irradiation. The pre-frontal area during the recovery phase elevated to the pre-irradiation phase

Based on our results, the cognitive and behavioral elements were different after exposure to EMFs from simulated high-pressure towers in the irradiated subjects. Behavioral analysis of experimental and control monkeys revealed that homogeneous behavioral indices, such as refreshment, sedentary, and natural appetite were observed in control monkeys, whereas experimental monkeys showed drowsiness, loss of appetite, sleep deprivation, fatigue, and depression after EMFs irradiation. These behavioral and cognitive changes in the tested monkey showed an increasing trend from the first day to the last day of the waving. On the other hand, from day 30 onwards, with the interruption of the waving, cognitive and behavioral changes returned to the initial state before the waving. It indicates the reversible effects of the fields caused by high-pressure rigs. According to research, people who live near high-pressure towers develop cognitive impairments (Aliyari, Hosseini, Menhaj, & Sahraei, 2019). The results of this study showed that exposure to high-frequency EMFs reduced melatonin secretion leading to sleep deprivation and immune deficiency in male Rhesus monkeys. sleep deprivation and immune deficiency are major contributors to depression (Burke, Davis, Otte, & Mohr, 2005; Hickie & Rogers, 2011; Thibodeau, Jorgensen, & Kim, 2006). Besides, high-frequency EMFs have been shown to reduce gene expression of the NMDA receptor and subsequently reduce plasma sodium and potassium levels. Sodium and potassium are the most important biological ions that cause irritability of the neurons and transmissions of nerve impulses. It is fundamental to note that sodium, potassium, and the NMDA receptor gene

are crucial elements in learning and memory processes (Basar, 2016; Clark & Mayer, 2016; De Zeeuw & Ten Brinke, 2015; Devesa, Lema, et al., 2016; Gruart, Leal-Campanario, López-Ramos, & Delgado-García, 2015). Exposure to extremely low-frequency EMFs (ELF-EMF 12Hz) has been shown to increase melatonin secretion. The metallothionein-1 (MT1) (or Mel1A) and metallothionein-2 (MT2) (or Mel1B) are two major receptors of melatonin which are protein-coupled receptors (GPCR). When receptors of melatonin bind to melatonin, the protein kinase A (PKA) or Protein kinase C (PKC) enzymes are activated (Kazemi et al., 2018a; Elahe Tekieh et al., 2018). Also, ELF-EMFs at a frequency of 12 Hz increase gene expression of N-methyl D-aspartate (NMDA), and subsequently improve visual memory of male Rhesus monkeys. The ELF EMFs stimulate glutamate on NMDA receptors and inhibits magnesium from the NMDA membrane receptor, resulting in more sodium outside and more potassium inside the cell (Kazemi et al., 2018a; Lau & Zukin, 2007; Salunke, Umathe, & Chavan, 2014). The highest density of the NMDA receptor gene is found in the hippocampus, amygdala, and prefrontal cortex in the central nervous system which plays a vital role in learning and memory processes (H Aliyari et al., 2018; Aliyari et al., 2015; Hamed Aliyari, Hedayat Sahraei, Mohammad Reza Daliri, Beh Minaei-Bidgoli, et al., 2018; Basar, 2016; Devesa, Lema, et al., 2016; Elahe Tekieh et al., 2018).

The findings of the current study also reveal that the plasma concentration of adrenaline increased after high-frequency EMF irradiation in the experimental monkeys. Adrenaline is a catecholamine secreted from the

central part of the adrenal gland, and when cortisol levels are low, adrenaline is secreted in response to anxiety and helps to chill and cope with tragedies. Increased adrenaline levels among study subjects may be due to impaired behaviors in this study. The activity of the adrenaline hormone is dependent on protein-coupled receptors (GPCR), and the activity of adenylate cyclase and increased cyclic adenosine monophosphate (cAMP) as a secondary messenger activates adrenaline (McGaugh & Roozendaal, 2002; Miao, Jänig, & Levine, 2000).

Hormones are chemicals released to develop and enhance cognitive capabilities, and any stimulus, environmental factor, or behavioral and cognitive impairments influence their secretion. Evidence shows that stress coping strategies in male Rhesus monkeys are different, and the cortisol and adrenaline secretion varies depending on the personal characteristics of the animals (Aliyari et al., 2015; Devesa, Almengló, & Devesa, 2016; Tekieh et al., 2017). Biochemical studies have shown that blood glucose increased after exposure to simulated high-frequency EMFs. Adrenaline (epinephrine) is shown to elevate blood glucose which is consistent with the findings of this study. Moreover, results from hematological studies indicated a reduction in RBC counts among irradiated monkeys after exposure to high-frequency EMFs. Reduced RBC counts (anemia) are associated with drowsiness, fatigue, immune deficiency, and depression in irradiated animals, which is consistent with the results of behavioral tests in subjects exposed to high-frequency EMFs radiation.

Previous studies have shown that exposure to high-voltage electric fields is associated with sleep and depression in male Rhesus monkeys (Hamed Aliyari, 2018; Hickie & Rogers, 2011). Results from hematological studies also showed an increase in the WBC counts after irradiation. Increased WBC counts as an infectious agent may impair normal immune function. The results from behavioral tests indicate a weakened immune system, drowsiness, and inactivity amongst subjects exposed to simulated high-frequency EMFs, which can be attributed to the immune deficiency of the male Rhesus monkey.

The anatomical assessment by MRI is a common and non-invasive technique used by neurologists to examine neurological disorders, such as Alzheimer's disease. Cerebral atrophy in MRI of the hippocampus indicates the presence of Alzheimer's disease. According to the results from anthropometric measurements of PFA by MRI via DICOM LiteBox, the irradiated subject showed

a decreasing trend in PFA after the irradiation compared to the control monkeys (Servien et al., 2008).

The frontal lobes are involved in behavior and personality. The smallest anatomical changes in the PFA can rapidly alter behavior and personality and cause behavioral and cognitive impairments. The frontal lobe is the vital part of the brain that plays a major role in cognition and behavior. Disruption in it disrupts natural behaviors and cognitive indicators (Kazemi et al., 2018a). Previous studies have shown that high-frequency EMFs can cause anatomical changes in the hippocampus and amygdala MRI of male Rhesus monkeys (Laakso et al., 1995; Soinen et al., 1994). Kazemi et al. argued that no changes were observed in hippocampus MRI of male Rhesus monkeys after exposure to low-frequency EMFs at a frequency of 12 Hz (Kazemi et al., 2018a).

It is essential to note that the recovery phase is a period in which the subject is recovered to the pre-irradiation level. Although the effect of high-frequency EMFs is temporary, the subject was irradiated 4h a day for four months. Besides, people living near high-voltage power lines are constantly exposed to high-frequency EMFs. Thus, further research is required to precisely examine the various pathological conditions.

5. Conclusion

The findings of the present study showed that the observable behavioral disorders, such as fatigue, anorexia, sleep deprivation, and depression in the experimental subject exposed to the electric field from simulated high-pressure towers were consistent with the results of other tests. The increase in adrenalin concentration increases glucose levels. Increased WBC counts cause immune deficiency. Decreased RBC counts lead to anemia, fatigue, and drowsiness in the experimental monkeys. All matched the monkey's behavior. It can be concluded that the EMFs from high-pressure electric towers are a serious hazard to cognitive abilities and can lead to mental disorders in humans and animals.

Ethical Considerations

Compliance with ethical guidelines

All ethical considerations based on the international standard were observed (IR.BUMS.REC.1394.112).

Funding

This research was conducted with the financial support of Baqiyatallah Neuroscience Research Center, Department of Electrical Engineering, [Amir Kabir University of Technology \(AUT\)](#) (Tehran Polytechnic) and [Qazvin Branch, Iran, Islamic Azad University](#) (research laboratory of intelligent systems and cognitive sciences)

Authors' contributions

Conceptualization: Hamed Aliyari, Seyed Hossein Hosseini, and Mohammad Bagher Menhaj; Methodology: Hamed Aliyari, Masoomeh Kazemi, Hedayat Sahraei, and Sahar Golabi; Investigation: Masoomeh Kazemi, Sahar Golabi, and Hamed Aliyari; Writing-original draft: Masoomeh Kazemi, Sahar Golabi, and Hamed Aliyari; Writing-review, and editing: Hedayat Sahraei, Hamed Aliyari, and Sahar Golabi; Resources: Seyed Hossein Hosseini; Supervision: Seyed Hossein Hosseini.

Conflict of interest

The authors declared no conflict of interest.

Acknowledgments

This Paper was extracted from a Ph.D. dissertation and funded by [Baqiyatallah University of Medical Sciences \(BMSU\)](#).

Reference

- Aliyari, H., Sahraei, H., Daliri, M. R., Minaei-Bidgoli, B., Kazemi, M., Agaei H, et al. (2018). The beneficial or harmful effects of computer game stress on cognitive functions of players. *Basic and Clinical Neuroscience*, 9(3):177. [DOI:10.29252/nirp.bcn.9.3.177]
- Aliyari, H., Sahraei, H., Erfani, M., Mohammadi, M., Kazemi, M., Daliri, M. R., et al. (2020). Changes in cognitive functions following violent and football video games in young male volunteers by studying brain waves. *BCN*, (3), 279-288. <https://bcn.iuums.ac.ir/article-1-1287-en.html>
- Aliyari, H., Sahraei, H., Golabi, S., Kazemi, M., Daliri, M. R., Minaei-Bidgoli, B. (2021). The effect of brain teaser games on the attention of players based on hormonal and brain signals changes. *Basic and Clinical Neuroscience*, 12(5), 587-596. [DOI:10.32598/bcn.2021.724.9]
- Aliyari, H., Kazemi, M., Tekieh, E., Salehi, M., Sahraei, H., & Daliri, MR, et al. (2015). The effects of Fifa 2015 computer games on changes in cognitive, hormonal and brain waves functions of young men volunteers. *Basic and Clinical Neuroscience*, 6(3), 193-201.
- Aliyari, H., Hosseini, S. H., Menhaj, M. B., & Sahraei, H. (2018) Analysis of the effects of high-voltage transmission line on human stress and attention through electroencephalography (EEG). *Iranian Journal of Science and Technology, Transactions of Electrical Engineering*, 2018, 1-8. [DOI:10.1007/s40998-018-0151-8]
- Aliyari, H., Hosseini, S., Sahraei, H., & Menhaj, M. (2019). Effect of proximity to high-voltage fields: Results of the neural network model and experimental model with macaques. *International Journal of Environmental Science and Technology*, 16(8), 4315-4326. [DOI:10.1007/s13762-018-1830-8]
- Aliyari, H., Sahraei, H., Erfanid, M., Tekieh, E., Salehib, M., Kazemib, M., et al. (2019). The impacts of video games on cognitive function and cortisol levels in young female volunteers. *Journal of Experimental and Clinical Neurosciences*, 6(1), 1-5. [DOI:10.13183/jecns.v6i1.87]
- Basar, E. (2016). Memory and brain dynamics: *Oscillations integrating attention, perception, learning, and memory*. CRC press. . [DOI:10.1201/9780203298732]
- Bayr, H. (2005). Reactive oxygen species. *Critical Care Medicine*, 33(12), S498-S501. [DOI:10.1097/01.CCM.0000186787.64500.12]
- Brady, T. F., Konkle, T., & Alvarez, G. A. (2011). A review of visual memory capacity: Beyond individual items and toward structured representations. *Journal of Vision*, 11(5), 4. [DOI:10.1167/11.5.4]
- Burke, H. M., Davis, M. C., Otte, C., & Mohr, D. C. (2005). Depression and cortisol responses to psychological stress: A meta-analysis. *Psychoneuroendocrinology*, 30(9), 846-856. [DOI:10.1016/j.psyneuen.2005.02.010]
- Clark, R. C., & Mayer, R. E. (2016). E-learning and the science of instruction: Proven guidelines for consumers and designers of multimedia learning. John Wiley & Sons. [DOI:10.1002/9781119239086]
- Cook, C., Thomas, A., & Prato, F. (2002). Human electrophysiological and cognitive effects of exposure to ELF magnetic and ELF modulated RF and microwave fields: A review of recent studies. *Bioelectromagnetics*, 23(2), 144-157. [DOI:10.1002/bem.107]
- De Pittà, M., Volman, V., Berry, H., & Ben-Jacob, E. (2011). A tale of two stories: A stroke regulation of synaptic depression and facilitation. *PLoS Computational Biology*, 7(12), e1002293. [DOI:10.1371/journal.pcbi.1002293]
- De Zeeuw, C. I., Ten Brinke, M. M. (2015). Motor learning and the cerebellum. *Cold Spring Harbor Perspectives in Biology*, 7(9), a021683. [DOI:10.1101/cshperspect.a021683]
- Devesa, J., Almengló, C., & Devesa, P. (2016). Multiple effects of growth hormone in the body: Is it really the hormone for growth? *Clinical Medicine Insights. Endocrinology and Diabetes*, 9, 47-71. [DOI:10.4137/CMED.S38201]

- Devesa, J., Lema, H., Zas, E., Munín, B., Taboada, P., & Devesa, P. (2016). Learning and memory recoveries in a young girl treated with growth hormone and neurorehabilitation. *Journal of clinical medicine*, 5(2), 14. [DOI:10.3390/jcm5020014]
- Duan, Y., Wang, Z., Zhang, H., He, Y., Fan, R., Cheng, Y., et al. (2014). Extremely low frequency electromagnetic field exposure causes cognitive impairment associated with alteration of the glutamate level, MAPK pathway activation and decreased CREB phosphorylation in mice hippocampus: Reversal by procyanidins extracted from the lotus seedpod. *Food & Function*, 5(9), 2289-2297. [DOI:10.1039/C4FO00250D]
- Fereidouni, A., Vahidi, B., Shishehgar, F., Mehr, T. H., & Tahmasbi, M. (2014). Human body modeling in the vicinity of high voltage transmission lines. *Science International*, 26(3). [Link]
- Garip, A., Akan, Z. (2010). Effect of ELF-EMF on number of apoptotic cells; correlation with reactive oxygen species and HSP. *Acta Biologica Hungarica*, 61(2), 158-167. [DOI:10.1556/ABiol.61.2010.2.4]
- Gruart, A., Leal-Campanario, R., López-Ramos, JC, & Delgado-García, J. M. (2015). Functional basis of associative learning and its relationships with long-term potentiation evoked in the involved neural circuits: Lessons from studies in behaving mammals. *Neurobiology of Learning and Memory*, 124, 3-18. [DOI:10.1016/j.nlm.2015.04.006]
- Harach, T., Marungruang, N., Dutilleul, N., Cheatham, V., McCoy, K., Neher, J., et al. (2015). Reduction of Alzheimer's disease beta-amyloid pathology in the absence of gut microbiota. arXiv preprint arXiv:150902273. [DOI:10.48550/arXiv.1509.02273]
- Hickie, I. B., & Rogers, N. L. (2011). Novel melatonin-based therapies: Potential advances in the treatment of major depression. *The Lancet*, 378(9791), 621-631. [DOI:10.1016/S0140-6736(11)60095-0]
- Kazemi, M., Sahraei, H., Aliyari, H., Tekieh, E., Saberi, M., Tavakoli, H., et al. (2018). Effect of extremely low frequency electromagnetic fields (ELF/MF) on NMDA-receptor gene expression and visual working memory of male rhesus macaques. *Basic and Clinical Neuroscience*, 9(3), 177-186. [DOI:10.29252/nirp.bcn.9.3.167]
- Laakso, M. P., Soininen, H., Partanen, K., Helkala, E. L., Hartikainen, P., Vainio, P., et al. (1995). Volumes of hippocampus, amygdala and frontal lobes in the MRI-based diagnosis of early Alzheimer's disease: correlation with memory functions. *Journal of Neural Transmission. Parkinson's Disease and Dementia Section*, 9(1), 73-86. [DOI:10.1007/BF02252964]
- LaBar, K. S., & Cabeza, R. (2006). Cognitive neuroscience of emotional memory. *Nature Reviews Neuroscience*, 7(1), 54-64. [DOI:10.1038/nrn1825]
- Lau, C. G., & Zukin, R. S. (2007). NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nature reviews. Neuroscience*, 8(6), 413-426. [DOI:10.1038/nrn2153]
- Liburdy, R., Sloma, T., Sokolic, R., Yaswen, P. (1993). ELF magnetic fields, breast cancer, and melatonin: 60 Hz fields block melatonin's oncostatic action on ER+ breast cancer cell proliferation. *Journal of Pineal Research*, 14(2), 89-97. [DOI:10.1111/j.1600-079X.1993.tb00491.x]
- Liburdy, R. P. (1992). Calcium signaling in lymphocytes and ELF fields evidence for an electric field metric and a site of interaction involving the calcium ion channel. *FEBS Letters*, 301(1), 53-59. [DOI:10.1016/0014-5793(92)80209-Y]
- McGaugh, J. L., & Roozendaal, B. (2002). Role of adrenal stress hormones in forming lasting memories in the brain. *Current Opinion in Neurobiology*, 12(2), 205-210. [DOI:10.1016/S0959-4388(02)00306-9]
- Miao, F. J., Janig, W., & Levine, J. D. (2000). Nociceptive neuroendocrine negative feedback control of neurogenic inflammation activated by capsaicin in the rat paw: Role of the adrenal medulla. *The Journal of Physiology*, 527(Pt 3), 601-610. [DOI:10.1111/j.1469-7793.2000.00601.x]
- Monti, M., Pernecco, L., Moruzzi, M., Battini, R., Zaniol, P., Barbiroli, B. (1991). Effect of ELF pulsed electromagnetic fields on protein kinase C activation process in HL-60 leukemia cells. *Journal of Bioelectricity*, 10(1-2), 119-130. [DOI:10.3109/15368379109031403]
- Murugappan, M., Ramachandran, N., & Sazali, Y. (2010). Classification of human emotion from EEG using discrete wavelet transform. *Journal of Biomedical Science and Engineering*, 3(04):390. [DOI:10.4236/jbise.2010.34054]
- Orban, G. A., Fize, D., Peuskens, H., Denys, K., Nelissen, K., Sanaert, S., et al. (2003). Similarities and differences in motion processing between the human and macaque brain: Evidence from fMRI. *Neuropsychologia*, 41(13), 1757-1768. [DOI:10.1016/S0028-3932(03)00177-5]
- Paz-Caballero, D., Cuertos, F., & Dobarro, A. (2006). Electrophysiological evidence for a natural/artifactual dissociation. *Brain Research*, 1067(1), 189-200. [DOI:10.1016/j.brainres.2005.10.046]
- Poiesz, B. J., Ruscetti, F. W., Gazdar, A. F., Bunn, P. A., Minna, J. D., Gallo, R. C. (1980). Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proceedings of the National Academy of Sciences*, 77(12), 7415-7419. [DOI:10.1073/pnas.77.12.7415]
- Salunke, B. P., Umathe, S. N., & Chavan, J. G. (2014). Involvement of NMDA receptor in low-frequency magnetic field-induced anxiety in mice. *Electromagnetic Biology and Medicine*, 33(4), 312-326. [DOI:10.3109/15368378.2013.839453]
- Servien, E., Viskontas, D., Giuffre, B. M., Coolican, M.R., & Parker, D. A. (2008). Reliability of bony landmarks for restoration of the joint line in revision knee arthroplasty. *Knee Surgery, Sports Traumatology, Arthroscopy*, 16(3), 263-269. [DOI:10.1007/s00167-007-0449-y]
- Sobel, E., & Davanipour, Z. (1996). Electromagnetic field exposure may cause increased production of amyloid beta and eventually lead to Alzheimer's disease. *Neurology*, 47(6), 1594-1600. [DOI:10.1212/WNL.47.6.1594]
- Soininen, H. S., Partanen, K., Pitkänen, A., Vainio, P., Hänninen, T., Hallikainen, M., et al. (1994). Volumetric MRI analysis of the amygdala and the hippocampus in subjects with age-associated memory impairment: Correlation to visual and verbal memory. *Neurology*, 44(9), 1660-1668. [DOI:10.1212/WNL.44.9.1660]
- Tekieh E, Riahi E, Kazemi M, Sahraei H, Tavakoli H, Aliyari H, et al. (2017). Role of basal stress hormones and amygdala dimensions in stress coping strategies of male rhesus mon-

keys in response to a hazard-reward conflict. *Iranian Journal of Basic Medical Sciences*, 20(8), 951-957. [DOI:10.22038/IJBMS.2017.9120]

Tekieh, E., Kazemi, M., Tavakoli, H., Saberi, M., Ghanaati, H., Hajinasrollah, M., et al. (2018). The effect of extremely low frequency electromagnetic fields on visual learning & memory and anatomical structures of the brain in male rhesus monkeys. *ISMJ*, 21(1), 40-53. [DOI:10.29252/ismj.21.1.40]

Thibodeau, R., Jorgensen, R. S., & Kim, S. (2006). Depression, anxiety, and resting frontal EEG asymmetry: A meta-analytic review. *Journal of Abnormal Psychology*, 115(4), 715-729. [DOI:10.1037/0021-843X.115.4.715]